

Asymptotic enumeration of sparse multigraphs with given degrees

Catherine Greenhill *

School of Mathematics and Statistics
The University of New South Wales
Sydney NSW 2052, Australia

csg@unsw.edu.au

Brendan D. McKay *

Research School of Computer Science
Australian National University
Canberra, ACT 0200, Australia

bdm@cs.anu.edu.au

15 March 2013

Abstract

Let J and J^* be subsets of $\mathbb{N}$ such that $0, 1 \in J$ and $0 \in J^*$. For infinitely many n , let $\mathbf{k} = (k_1, \dots, k_n)$ be a vector of nonnegative integers whose sum M is even. We find an asymptotic expression for the number of multigraphs on the vertex set $\{1, \dots, n\}$ with degree sequence given by $\mathbf{k}$, such that every loop has multiplicity in J^* and every non-loop edge has multiplicity in J . Equivalently, these are symmetric integer matrices with values J^* allowed on the diagonal and J off the diagonal. Our expression holds when the maximum degree $k_{\max}$ satisfies $k_{\max} = o(M^{1/3})$. We prove this result using the switching method, building on an asymptotic enumeration of simple graphs with given degrees (McKay and Wormald, 1991). Our application of the switching method introduces a novel way of combining several different switching operations into a single computation.

1 Introduction

Let $k_{i,n}$ be a nonnegative integer for all pairs (i, n) of integers which satisfy $1 \leq i \leq n$. Then for each $n \geq 1$, let $\mathbf{k} = \mathbf{k}(n) = (k_{1,n}, \dots, k_{n,n})$. We usually write k_i instead of $k_{i,n}$. Define

*Research supported by the Australian Research Council.

$M = \sum_{i=1}^n k_i$. We assume that M is even for an infinite number of values of n , and tacitly restrict ourselves to such n .

For subsets J, J^* of $\mathbb{N}$, define $\mathcal{G}(\mathbf{k}, J, J^*)$ to be the set of all multigraphs on the vertex set $\{1, \dots, n\}$ with degree sequence given by $\mathbf{k}$ such that the multiplicity of every loop belongs to J^* and the multiplicity of every non-loop edge belongs to J . Loops contribute 2 to the degree of their vertex. Also define $\mathcal{G}(\mathbf{k})$ to be the set of multigraphs on $\{1, \dots, n\}$ with degree sequence $\mathbf{k}$ (and no restrictions on the multiplicities).

Equivalently we may think of $\mathcal{G}(\mathbf{k}, J, J^*)$ as the set of all $n \times n$ symmetric matrices $A = (a_{ij})$ with all diagonal entries in J^* , all off-diagonal entries in J and with

$$2a_{ii} + \sum_{j \neq i} a_{ij} = k_i \quad (1.1)$$

for $i = 1, 2, \dots, n$. Note that diagonal entries are weighted by two in the row sum.

Let $k_{\max} = \max_{i=1}^n k_i$. We find an asymptotic expression for

$$G(\mathbf{k}, J, J^*) = |\mathcal{G}(\mathbf{k}, J, J^*)|$$

that holds when $k_{\max} = o(M^{1/3})$. For $r = 1, 2, \dots$ let

$$M_r = \sum_{i=1}^n [k_i]_r$$

where $[a]_r = a(a-1)(a-2) \cdots (a-r+1)$ denotes the falling factorial. Then $M_1 = M$ and $M_r \leq k_{\max} M_{r-1}$ for all $r \geq 2$.

For $i \geq 0$, define

$$x_i = \begin{cases} 1, & \text{if } i \in J, \\ 0, & \text{otherwise;} \end{cases} \quad y_i = \begin{cases} 1, & \text{if } i \in J^*, \\ 0, & \text{otherwise.} \end{cases}$$

Our main result is the following.

Theorem 1.1. *Let J and J^* be subsets of $\mathbb{N}$ such that $0, 1 \in J$ and $0 \in J^*$. Suppose that $n \rightarrow \infty$, $M \rightarrow \infty$ and $k_{\max}^3 = o(M)$. Then*

$$\begin{aligned} G(\mathbf{k}, J, J^*) &= \frac{M!}{(M/2)! 2^{M/2} k_1! \cdots k_n!} \exp \left(\left(y_1 - \frac{1}{2} \right) \frac{M_2}{M} + \left(x_2 - \frac{1}{2} \right) \frac{M_2^2}{2M^2} + \frac{M_2^4}{4M^5} \right. \\ &\quad \left. - \frac{M_2^2 M_3}{2M^4} + \left(x_3 - x_2 + \frac{1}{3} \right) \frac{M_3^2}{2M^3} + O(k_{\max}^3/M) \right). \end{aligned}$$

For convenience, we restate the formula in the regular case.

Corollary 1.2. *Suppose that $\mathbf{k} = (k, k, \dots, k)$ with kn even, such that $k = o(n^{1/2})$ as $n \rightarrow \infty$. Then*

$$G(\mathbf{k}, J, J^*) = \frac{(kn)!}{(kn/2)! 2^{kn/2} (k!)^n} \exp(-Q(k, n) + O(k^2/n))$$

where

$$Q(k, n) = \frac{1}{4} (k-1)((-1)^{x_2}(k-1) + 2(-1)^{y_1}) + \frac{k^3}{12n}(6x_2 - 6x_3 + 1).$$

Setting $y_1 = x_2 = x_3 = 0$ we recapture the asymptotic expression for sparse simple graphs with given degrees presented in [12, Theorem 5.2]. Similarly, setting $y_1 = 1, x_2 = x_3 = 0$ we obtain the asymptotic enumeration by degree sequence of sparse graphs with loops allowed but no multiple edges: this is the second expression in [4, Theorem 1.5].

We remark that the conditions $0 \in J^*$ and $0, 1 \in J$ in Theorem 1.1 can be replaced by weaker conditions: namely, that J^* is non-empty and that J has at least two elements with the smallest two consecutive. Let s denote the least element of J^* and let $t, t+1$ be the smallest two elements of J . This case is reduced to ours by subtracting s from each diagonal element, t from each off-diagonal element, and $2s + (n-1)t$ from each k_i , provided the new degree sequence satisfies the conditions of Theorem 1.1.

Theorem 1.1 is proved using the switching method, building on an asymptotic enumeration of simple graphs with given degrees [12]. Our application of the switching method introduces a novel way of combining several different switching operations into a single computation.

The remainder of the paper is structured as follows. The history of this asymptotic enumeration problem is briefly reviewed in Section 1.1, then the new switching theorem is given in Section 2. In Section 3 we describe 15 types of switchings on multigraphs, which are used to show that certain multiplicities of edges or loops are rare. These switchings are analysed together in Subsection 3.14 using the new switching theorem. In Section 4 we complete the enumeration with the help of some calculations performed in [12].

Finally, in Section 5 we show that a naïve argument leads to a formula for $G(\mathbf{k}, J, J^*)$ which differs asymptotically by a constant factor from the result of Theorem 1.1 in the regular case. The constant factor takes two different values, depending on whether $2 \in J$.

1.1 History

The earliest work on this problem was that of Read [14], who found exact and asymptotic formulae when $k_1 = \dots = k_n = 3$, for all four combinations of $J^* \in \{\{0\}, \mathbb{N}\}$ and $J \in$

$\{\{0, 1\}, \mathbb{N}\}$. The best result for $J^* = \{0\}, J = \{0, 1\}$ in the sparse range is that of McKay and Wormald [12] who treated $k_{\max} = o(M^{1/3})$; see that paper for a survey of the many earlier results on the 0-1 case. In addition to $J^* = \{0\}, J = \{0, 1\}$, Bender and Canfield [2] found the asymptotics for $J^* = \{0\}, J = \mathbb{N}$ when $k_{\max} = O(1)$. (Although that paper allows nonzero diagonal entries, they contribute singly to the row sums, not doubly as we have it.)

In [11], McKay and Wormald considered $J^* = \{0\}, J = \{0, 1\}$ in the dense domain defined by $\min\{k, n - k - 1\} \geq cn/\log n$ for $c > \frac{2}{3}$ and $|k_i - k| = O(n^{1/2+\varepsilon})$ for all i , where k is the average degree. McKay [8] found a better error term under the same conditions, while Barvinok and Hartigan allowed a wider range of degrees [1]. Greenhill and McKay [4] added the option of loops in both the sparse and dense ranges by considering $J^* = \{0, 1\}, J = \{0, 1\}$. Finally, McLeod and McKay [9] analysed the case of $J^* = \{0\}, J = \mathbb{N}$ when $k_1 = \dots = k_n > cn/\log n$ for $c > \frac{1}{6}$.

For sparse rectangular matrices which are not necessarily symmetric, see Greenhill and McKay [3].

2 The switching theorem

In order to bound the number of multigraphs having some unusual properties, we will apply the method of switchings. It will be necessary to apply several switching types, which we could analyse one at a time as in [3]. Instead we now prove a generalized switching theorem that allows us to analyse all of them at once.

Theorem 2.1. *Let $\Gamma = \Gamma(V, E)$ be a directed multigraph, and let $\alpha : E \rightarrow \mathbb{R}_+$ be a positive weighting of the edges of Γ . Fix a non-empty finite set C , whose elements we will call colours, and let $c : E \rightarrow C$ be an edge colouring of Γ (which need not be proper). For all $v \in V$ and $c \in C$ denote by $\Gamma_c^-(v)$ the set of edges of colour c entering v , and by $\Gamma_c^+(v)$ the set of edges of colour c leaving v .*

We introduce the set of variables $\{N(v) : v \in V\} \cup \{s(e) : e \in E\}$ and consider the following system of linear inequalities on these variables:

$$N(v) \geq 0, \quad (v \in V) \tag{2.1}$$

$$s(e) \geq 0, \quad (e \in E) \tag{2.2}$$

$$\sum_{e \in \Gamma_c^-(v)} s(e) \leq N(v), \quad (v \in V, c \in C) \tag{2.3}$$

$$\sum_{e \in \Gamma_c^+(v)} \alpha(e)s(e) \geq N(v), \quad (v \in V, c \in C, \Gamma_c^+(v) \neq \emptyset). \tag{2.4}$$

Let $C(v) = \{c \in C : \Gamma_c^-(v) \neq \emptyset\}$ be the set of colours entering v . For each $v \in V$ which is not a source, let $\lambda_c(v), c \in C(v)$ be positive numbers such that $\sum_{c \in C(v)} \lambda_c(v) \leq 1$. For each $vw \in E$, define $\hat{\alpha}(vw) = \alpha(vw)/\lambda_{c(vw)}(w)$. Extend this function to any directed path P by defining $\hat{\alpha}(P) = \prod_{e \in P} \hat{\alpha}(e)$.

Now suppose that $X, Y, Z \subseteq V$ satisfy the following conditions:

1. $Z \neq \emptyset$, $Z \subseteq X$, and $Y \cap Z = \emptyset$;
2. If $v \in V$ is a sink of Γ , or if $\hat{\alpha}(vw) \geq 1$ for some $vw \in E$, then $v \in Z$.

For any $W, W' \subseteq V$, define $\mathcal{P}(W, W')$ to be the set of non-trivial directed paths in Γ which start in W , end in W' , and have no internal vertices in $Y \cup Z$. Then every solution of (2.1)–(2.4) satisfies

$$\sum_{v \in Y} N(v) \leq \frac{\max_{P \in \mathcal{P}(Y, Z)} \hat{\alpha}(P)}{1 - \max_{P \in \mathcal{P}(Y, Y)} \hat{\alpha}(P)} \sum_{v \in X} N(v), \quad (2.5)$$

where maximums over empty sets are taken to be 0.

Proof. The case of one colour appears in [6, Theorem 3], apart from an inconsequential difference in the conditions on Z . We proceed by reducing the general problem to one in which there is only one colour, then modifying Z slightly so that [6, Theorem 3] applies.

Define $\Gamma^-(v)$ and $\Gamma^+(v)$ to be the set of all edges (regardless of colour) entering v or leaving v , respectively. For each edge $vw \in E$, define $\hat{s}(vw) = s(vw)\lambda_{c(vw)}(w)$. If we weight inequality (2.3) by $\lambda_c(v)$ and sum over $c \in C(v)$, we obtain

$$\sum_{e \in \Gamma^-(v)} \hat{s}(e) \leq N(v), \quad (v \in V). \quad (2.6)$$

Similarly, if $\Gamma^+(v) \neq \emptyset$, then we can sum (2.4) over those $c \in C$ with $\Gamma_c^+(v) \neq \emptyset$ to obtain

$$\sum_{e \in \Gamma^+(v)} \hat{\alpha}(e)\hat{s}(e) \geq N(v), \quad (v \in V, \Gamma^+(v) \neq \emptyset). \quad (2.7)$$

Together with the nonnegativity of $N(v)$ and $\hat{s}(e)$, we have equations of the form of (2.1)–(2.4) with only one colour.

Next remove from Γ all edges vw where $v \in Z$. Doing so can only weaken the conditions, by decreasing the left hand side of some inequalities in (2.6) or removing some inequalities in (2.7). Note that none of the quantities in (2.5) are changed by removal of these edges. Now Z satisfies the requirements of [6, Theorem 3], and applying that theorem to the system defined by (2.6), (2.7) and the nonnegativity of $N(v)$ and $\hat{s}(e)$ gives (2.5) immediately. $\square$

We now describe how Theorem 2.1 can be used for counting.

Suppose we have a finite set of “objects” partitioned into disjoint classes $\mathcal{S}(v)$, where $v \in V$ for some index set V . Define $N(v) = |\mathcal{S}(v)|$ for each $v \in V$. Also suppose that for each $c \in C$ we have a relation Ψ_c between objects: to be precise, Ψ_c is a multiset of ordered pairs (Q, R) of objects. (We call Ψ_c a *switching* and usually define it by some operation that modifies Q to make R .)

Now define an edge-coloured directed multigraph $\Gamma = (V, E)$ with vertex set V , where Γ has a directed edge vw of colour c if and only if $(Q, R) \in \Psi_c$ for some $Q \in \mathcal{S}(v)$ and $R \in \mathcal{S}(w)$. (There is at most one edge of each colour between any pair of distinct vertices of Γ .) For each $vw \in E$ let $s'(vw) = |\{(Q, R) \in \Psi_c : Q \in \mathcal{S}(v), R \in \mathcal{S}(w)\}|$, counting multiplicities, where c is the colour of vw .

Fix $v \in V$ and $c \in C$ such that $\Gamma_c^+(v) \neq \emptyset$. Suppose that for any $Q \in \mathcal{S}(v)$ there are at least $a_c(v) > 0$ objects R with $(Q, R) \in \Psi_c$, counting multiplicities. Then

$$\sum_{e \in \Gamma_c^+(v)} s'(e) \geq a_c(v)N(v).$$

Similarly, for fixed $v \in V$ and $c \in C$, suppose that for every $R \in \mathcal{S}(w)$ there are at most $b_c(w) > 0$ objects Q with $(Q, R) \in \Psi_c$, counting multiplicities. Then

$$\sum_{e \in \Gamma_c^-(w)} s'(e) \leq b_c(w)N(w).$$

Defining $s(vw) = s'(vw)/b_{c(vw)}(w)$ and $\alpha(vw) = b_{c(vw)}(w)/a_{c(vw)}(v)$ we obtain equations (2.1)–(2.4). Theorem 2.1 can thus be used to bound the relative values of $\sum_{v \in X} |\mathcal{S}(v)|$ and $\sum_{v \in Y} |\mathcal{S}(v)|$ if there is some Z for which X, Y, Z satisfy the requirements of the lemma. A common choice (which we will use here) is $X = V$, as then we bound the fraction of all objects which lie in $\bigcup_{v \in Y} \mathcal{S}(v)$.

3 Switchings on multigraphs

Define

$$\begin{aligned} N_1 &= \max\{\lceil \log M \rceil, \lceil 480M_2/M \rceil\}, \\ N_2 &= \max\{\lceil \log M \rceil, \lceil 240M_2^2/M^2 \rceil\}, \\ N_3 &= \max\{\lceil \log M \rceil, \lceil 240M_3^2/M^3 \rceil\}. \end{aligned} \tag{3.1}$$

(We have not attempted to optimise constants.) Given a multigraph Q , let $\ell_D(Q)$ denote the number of loops with multiplicity D and let $e_D(Q)$ denote the number of non-loop edges with multiplicity D , for $D \geq 1$.

Let

$$\mathcal{G}_0 = \mathcal{G}(\mathbf{k}, J \cup \{4, 5, 6, \dots\}, J^* \cup \{2, 3, 4, \dots\})$$

be the set of all multigraphs with degree sequence $\mathbf{k}$, and allowing all multiplicities except for those in $\{1\} - J^*$ on loops and those in $\{2, 3\} - J$ on non-loops. Note that $\mathcal{G}(\mathbf{k}, J, J^*) \subseteq \mathcal{G}_0$.

We also define the subsets Y, Z of $\mathcal{G}_0$ by

$$Y = \mathcal{G}_0 - \{Q \in \mathcal{G}_0 \mid \ell_D(Q) = 0 \text{ for } D \geq 2, e_D(Q) = 0 \text{ for } D \geq 4, \\ \ell_1(Q) \leq N_1, e_2(Q) \leq N_2, e_3(Q) \leq N_3\}, \quad (3.2)$$

$$Z = \{Q \in \mathcal{G}_0 \mid \ell_D(Q) = 0 \text{ for } D \geq 2, e_D(Q) = 0 \text{ for } D \geq 4, \\ \ell_1(Q) \leq \lceil N_1/2 \rceil, e_2(Q) \leq \lceil N_2/2 \rceil, e_3(Q) \leq \lceil N_3/2 \rceil\}. \quad (3.3)$$

We define 15 coloured switchings which act to reduce the number of loops or edges with high multiplicities, moving in steps from Y towards Z . These switchings are defined below, together with a description of when each should be used. Indeed, any given switching will only be used on multigraphs in $\mathcal{G}_0$ for which none of the switchings with a lower-labelled colour are applicable. An important property of all the switchings is that they do not create simple loops (with multiplicity 1) and they do not create non-loop edges of multiplicity 2 or 3, as these may not be allowed for multigraphs in $\mathcal{G}_0$.

For each switching colour c and multigraphs Q, R we will define the following parameters:

- $a_c(Q)$ is a lower bound on the number of ways in which a switching of colour c can be applied to Q ;
- $b_c(R)$ is an upper bound on the number of ways in which a switching of colour c can produce R ;
- if R can be obtained from Q by performing a switching of colour c , then we let $\alpha(Q, R)$ be an upper bound on $b_c(R)/a_c(Q)$. (Note that the colour c is determined by the edge QR .)

Finally in Section 3.14 we will analyse all the switchings at once by applying Theorem 2.1.

3.1 Switchings of colour 1

A switching of colour 1 is used to reduce the number of loops of multiplicity equal to 2 or multiplicity at least 4. It is applied to multigraphs $Q \in \mathcal{G}_0$ with $L(Q) > \lceil 3M^{1/2} \rceil$, where

$$L(Q) = \ell_2(Q) + \sum_{D \geq 4} \ell_D(Q) \quad (3.4)$$

is the number of loops in Q with multiplicity 2 or multiplicity at least 4. The switching is described by the sequence (v_1, v_2) of distinct vertices such that there is a loop of multiplicity D_1 at v_1 and a loop of multiplicity D_2 at v_2 , with $D_1, D_2 \in \{2\} \cup \{4, 5, \dots\}$.

Let m be the multiplicity of the edge from v_1 to v_2 in Q (which may equal zero). The switching reduces the multiplicity of both loops by 2, and increases the multiplicity of the edge $\{v_1, v_2\}$ by 4. This operation is depicted in Figure 1.




Figure 1: A switching of colour 1

Suppose that R can be produced from Q using a switching of colour 1. The number of ways to perform a switching of colour 1 in Q is exactly

$$[L(Q)]_2 \geq 9M$$

as there is no restriction on the value of m . The number of ways that R can be produced using a switching of colour 1 is bounded above by $M/4$, which is a bound on the number of ways to choose an oriented edge with multiplicity at least 4. Hence we can set $a_1(Q) = 9M$ and $b_1(R) = M/4$, giving

$$\alpha(Q, R) = \frac{1}{36}.$$

3.2 Switchings of colour 2

A switching of colour 2 is used to reduce the number of loops of multiplicity three to below $\lceil 3M^{1/2} \rceil$. It is performed for multigraphs Q for which switchings of colour 1 do not apply and such that $\ell_3(Q) > \lceil 3M^{1/2} \rceil$. The switching is described by the sequence (v_1, v_2) of distinct vertices such that there is a loop of multiplicity 3 at v_1 and at v_2 .

Let m be the multiplicity of the edge from v_1 to v_2 in Q (which may equal zero). The switching removes the loop at v_1 and at v_2 , and increases the multiplicity of the edge $\{v_1, v_2\}$ by 6, as illustrated in Figure 2.

Suppose that the multigraph R can be produced from Q using a switching of colour 2. The number of ways to perform a switching of colour 2 in Q is exactly $[\ell_3(Q)]_2 \geq 9M$, and the number of ways that R can be produced using a switching of colour 2 is at most $M/6$




Figure 2: A switching of colour 2

(since an oriented edge of multiplicity at least 6 determines the reverse operation, from R to Q). Hence we can take $a_2(Q) = 9M$ and $b_2(R) = M/6$, leading to

$$\alpha(Q, R) = \frac{1}{54}.$$

3.3 Switchings of colour 3

A switching of colour 3 is used to reduce the number of loops of multiplicity 1 to below $\lceil M^{1/2} \rceil$. It is applied to multigraphs Q for which switchings of colour 1 and 2 do not apply and such that $\ell_1(Q) > \lceil M^{1/2} \rceil$. The switching is defined by the sequence (v_1, v_2, v_3) of distinct vertices such that each of v_1, v_2, v_3 has a simple loop in Q (of multiplicity 1), and none of the edges $\{v_1, v_2\}, \{v_1, v_3\}, \{v_2, v_3\}$ are present in Q . The switching removes these three loops and joins the three vertices pairwise by simple edges (of multiplicity 1), as illustrated in Figure 3.




Figure 3: A switching of colour 3

Suppose that the multigraph R can be produced from Q using a switching of colour 3. The number of ways to perform a switching of colour 3 in Q is at least

$$[\ell_1(Q)]_3 - O(k_{\max} \ell_1(Q)^2) = \ell_1(Q)^3(1 - o(1)) \geq \frac{1}{2} M^{3/2},$$

while the number of ways that R can be produced using a switching of colour 3 is at most

$k_{\max}M$. Therefore we can take $a_3(Q) = \frac{1}{2}M^{3/2}$ and $b_3(R) = k_{\max}M$, leading to

$$\alpha(Q, R) = \frac{2k_{\max}}{M^{1/2}} = o(1).$$

3.4 Switchings of colour 4

A switching of colour 4 is used to reduce the number of non-loop edges of multiplicity greater than $\max\{4, \lceil k_{\max}^{1/2} \rceil\}$. It is applied to multigraphs Q for which switchings of colours 1, 2, 3 do not apply and such that $E^+(Q) > \lceil 4k_{\max}^{1/2} M^{1/2} \rceil$, where

$$E^+(Q) = \sum_{D=\max\{4, \lceil k_{\max}^{1/2} \rceil\}+1}^{k_{\max}} e_D(Q). \quad (3.5)$$

A switching of colour 4 in Q is described by a sequence (v_1, w_1, v_2, w_2) of distinct vertices such that

- the multiplicity of edge (v_1, w_1) in Q is D_1 and the multiplicity of (v_2, w_2) in Q is D_2 , where $D_1, D_2 > \max\{4, \lceil k_{\max}^{1/2} \rceil\}$,
- the edges $\{v_1, v_2\}$ and $\{w_1, w_2\}$ have multiplicity zero in Q (they are non-edges).

The switching reduces the multiplicity of edges (v_1, w_1) and (v_2, w_2) by one and gives multiplicity 1 to edges $\{v_1, v_2\}$ and $\{w_1, w_2\}$, as shown in Figure 4.




Figure 4: A switching of colour 4 or 5

Now suppose that the multigraph R can be produced from Q using a switching of colour 4. The number of switchings of colour 4 that can be performed in Q is bounded below by

$$4[E^+(Q)]_2 - O(k_{\max}^2 E^+(Q)) = 4E^+(Q)^2(1 - o(1)) \geq 64k_{\max} M(1 - o(1)) > 60k_{\max} M,$$

and the number of switchings of colour 4 that can produce R is at most

$$\frac{2k_{\max}^2 M}{\max\{4, \lceil k_{\max}^{1/2} \rceil\}^2} \leq 2k_{\max} M.$$

Hence we can take $a_4(Q) = 60 k_{\max} M$ and $b_4(R) = 2k_{\max} M$, leading to

$$\alpha(Q, R) = \frac{1}{30}.$$

3.5 Switchings of colour 5

A switching of colour 5 is used to reduce the number of non-loop edges of multiplicity at least 5 and at most $\lceil k_{\max}^{1/2} \rceil$. For a multigraph $Q \in \mathcal{G}_0$, let

$$E^-(Q) = \sum_{D=5}^{\lceil k_{\max}^{1/2} \rceil} e_D(Q).$$

Switchings of colour 5 are applied to multigraphs Q for which switchings of colours $1, \dots, 4$ do not apply and such that $E^-(Q) > \lceil 3k_{\max} M^{1/2} \rceil$.

A switching of colour 5 is described by a sequence (v_1, w_1, v_2, w_2) of distinct vertices such that

- the multiplicity of $\{v_1, w_1\}$ in Q is D_1 and the multiplicity of $\{v_2, w_2\}$ in Q is D_2 , where

$$D_1, D_2 \in \{5, 6, \dots, \lceil k_{\max}^{1/2} \rceil\},$$

- the multiplicity of $\{v_1, v_2\}$ and $\{w_1, w_2\}$ in Q is zero (these are non-edges).

This switching is also illustrated by Figure 4, but with different conditions on D_1, D_2 as above.

Suppose that R can be produced from Q using a switching of colour 5. The number of ways that a switching of colour 5 can be performed in Q is bounded below by

$$4[E^-(Q)]_2 - O(k_{\max}^2 E^-(Q)) = 4E^-(Q)^2(1 - o(1)) \geq 30k_{\max}^2 M,$$

and the number of ways to produce R using a switching of colour 5 is at most

$$k_{\max}^2 M.$$

Hence we can let $a_5(Q) = 30k_{\max}^2 M$ and $b_5(R) = k_{\max}^2 M$, leading to

$$\alpha(Q, R) = \frac{1}{30}.$$

3.6 Switchings of colour 6, 7, 8

A switching of colour $4 + j$ is used to reduce the number of edges of multiplicity j to below $\lceil M^{5/6} \rceil$, for $j = 2, 3, 4$. Switchings of colour $4 + j$ are applied to multigraphs Q for which switchings of colours $1, \dots, 3 + j$ do not apply and such that

$$e_j(Q) > \lceil M^{5/6} \rceil.$$

Given such a multigraph Q , a switching of colour $4 + j$ is defined by a sequence

$$(v_1, w_1, v_2, w_2, \dots, v_j, w_j)$$

of distinct vertices such that

- (v_r, w_r) is an edge of multiplicity j in Q for $r = 1, \dots, j$,
- every edge $\{v_r, w_s\}$ with $1 \leq r \neq s \leq j$ has multiplicity 0 in Q (it is a non-edge).

The switching deletes these j edges of multiplicity j , and inserts a complete bipartite graph $K_{j,j}$ on $\{v_1, \dots, v_j\} \cup \{w_1, \dots, w_j\}$, with all j^2 new edges simple (that is, multiplicity 1). This operation is illustrated in the Figure 5 for the case $j = 4$.




Figure 5: A switching of colour 8

Suppose that the multigraph R can be obtained from Q using a switching of colour $4 + j$, for $j \in \{2, 3, 4\}$. The number of ways in which a switching of colour $4 + j$ can be performed in Q is bounded below by

$$2^j [e_j(Q)]_j - O(k_{\max}^2 e_j(Q)^{j-1}) \geq \frac{1}{2} M^{5j/6},$$

while the number of switchings of colour $4 + j$ which produce R is at most $k_{\max}^{2j-2} M$. Hence we can define $a_{4+j}(Q) = \frac{1}{2} M^{5j/6}$ and $b_{4+j}(R) = k_{\max}^{2j-2} M$ for $j \in \{2, 3, 4\}$. Since $j \geq 2$, this leads to

$$\alpha(Q, R) = \frac{2k_{\max}^{2j-2} M}{M^{5j/6}} = O\left(\frac{k_{\max}^2}{M^{2/3}}\right) = o(1).$$

Before describing more coloured switchings, we prove a useful fact.

Lemma 3.1. *Suppose that $Q \in \mathcal{G}_0$ is such that no switching of colour 1 to 8 applies to Q . Then $e_1(Q) = (\frac{1}{2} - o(1))M$.*

Proof. Since no switching of colour 1, 2 or 3 applies we have

$$\sum_{D \geq 1} D \ell_D(Q) = O(k_{\max} M^{1/2}) = O(M^{5/6}),$$

and since no switching of colour 4–8 applies we know that

$$\begin{aligned} \sum_{D \geq 2} D e_D(Q) &\leq 3 \lceil M^{5/6} \rceil + k_{\max} E^+(Q) + \lceil k_{\max}^{1/2} \rceil E^-(Q) \\ &= O(M^{5/6} + k_{\max}^{3/2} M^{1/2}) \\ &= o(M). \end{aligned}$$

The result follows. □

3.7 Switchings of colour 9

Switchings of colour 9 reduce the number of loops of multiplicity 2 or multiplicity at least 4, until this number is zero. They are applied to multigraphs Q for which the switchings of colours 1, $\dots$, 8 do not apply and such that $L(Q) \geq 1$.

Let Q be such a multigraph. A switching of colour 9 in Q is defined by a sequence $(v_0, v_1, w_1, v_2, w_2)$ of distinct vertices such that

- there is a loop at vertex v_0 in Q with multiplicity D , where $D = 2$ or $D \geq 4$,
- $\{v_1, w_1\}$ and $\{v_2, w_2\}$ are simple edges (of multiplicity 1),
- $\{v_0, v_1\}, \{v_0, v_2\}, \{v_0, w_1\}, \{v_0, w_2\}$ are all non-edges in Q (with multiplicity zero).

The switching reduces the multiplicity of the loop at v_0 to $D - 2$, removes the two simple edges $\{v_1, w_1\}$ and $\{v_2, w_2\}$ and inserts the four simple edges $\{v_0, v_1\}, \{v_0, v_2\}, \{v_0, w_1\}, \{v_0, w_2\}$, as shown in Figure 6.

Suppose that the multigraph R can be produced from Q by a switching of colour 9. The number of ways to perform a switching of colour 9 in Q is at least

$$L(Q) \left([2e_1(Q)]_2 - O(k_{\max}^2 e_1(Q)) \right) \geq \frac{1}{2} L(Q) M^2,$$




Figure 6: A switching of colour 9

using Lemma 3.1, while the number of ways to produce R using a switching of colour 9 is at most M_4 . (We ignore the presence of the loop at v_0 in this upper bound, since no such loop exists when $D = 2$.)

Hence we can let $a_9(Q) = \frac{1}{2}L(Q)M^2$ and $b_9(R) = M_4$, leading to

$$\alpha(Q, R) = \frac{2M_4}{L(Q)M^2} = O\left(\frac{k_{\max}^3}{M}\right)$$

since $L(Q) \geq 1$.

3.8 Switchings of colour 10

Switchings of colour 10 reduce the number of loops of multiplicity 3, until this number is zero. They are applied to multigraphs Q such that switchings of colours $1, \dots, 9$ do not apply and such that $\ell_3(Q) \geq 1$.

Let Q be such a multigraph. A switching of colour 10 is defined by a sequence of distinct vertices $(v_0, v_1, w_1, v_2, w_2, v_3, w_3)$ such that

- there is a loop of multiplicity 3 at v_0 in Q ,
- there is a simple edge $\{v_j, w_j\}$ in Q , for $j = 1, 2, 3$,
- the edges $\{v_0, v_j\}$ and $\{v_0, w_j\}$ all have multiplicity 0 in Q , for $j = 1, 2, 3$.

The switching removes the loop of multiplicity 3 and the simple edges $\{v_j, w_j\}$ for $j = 1, 2, 3$, and adds the six simple edges $\{\{v_0, v_j\}, \{v_0, w_j\} \mid j = 1, 2, 3\}$, as shown in Figure 7.

Now let R be a multigraph which can be formed from Q by a switching of colour 10. The number of ways that a switching of colour 10 can be performed in Q is at least

$$\ell_3(Q) ([2e_1(Q)]_3 - O(k_{\max}^2 e_1(Q)^2)) \geq \frac{1}{2} \ell_3(Q) M^3,$$




Figure 7: A switching of colour 10

using Lemma 3.1, and the number of ways that R can be produced using a switching of colour 10 is bounded above by M_6 . Hence we can take $a_{10}(Q) = \frac{1}{2} \ell_3(Q) M^3$ and $b_{10}(R) = M_6$. This leads to

$$\alpha(Q, R) = \frac{2 M_6}{\ell_3(Q) M^3} = O\left(\frac{k_{\max}^5}{M^2}\right) = o\left(\frac{k_{\max}^3}{M}\right).$$

We do not tackle single loops yet. Instead, the next two switchings reduce non-loop edges of high multiplicity down to zero.

3.9 Switchings of colour 11

Switchings of colour 11 reduce the number of non-loop edges with multiplicity 4 or multiplicity at least 7, until this number is zero. Switchings of colour 11 are applied to multigraphs Q for which switchings of colours $1, \dots, 10$ do not apply and $E(Q) \geq 1$, where

$$E(Q) = e_4(Q) + \sum_{D \geq 7} e_D(Q).$$

Let Q be such a multigraph. A switching of colour 11 in Q is defined by a sequence

$$(v_0, w_0, v_1, w_1, v_2, w_2, v_3, w_3)$$

of distinct vertices such that

- the edge $\{v_0, w_0\}$ has multiplicity D in Q , where $D = 4$ or $D \geq 7$,
- edges $\{v_j, w_j\}$ are simple edges in Q (with multiplicity 1) for $j = 1, 2, 3$,
- the edges $\{v_0, v_j\}$ and $\{w_0, w_j\}$ are non-edges in Q for $j = 1, 2, 3$ (with multiplicity zero).

The switching reduces the multiplicity of $\{v_0, w_0\}$ to $D - 3$, removes the edges $\{v_j, w_j\}$ for $j = 1, 2, 3$ and adds the simple edges $\{v_0, v_j\}$ and $\{w_0, w_j\}$ for $j = 1, 2, 3$. This operation is illustrated in Figure 8.




Figure 8: A switching of colour 11

Suppose that the multigraph R can be obtained from Q by performing a switching of colour 11. The number of ways to perform a switching of colour 11 in Q is at least

$$2E(Q) ([2e_1(Q)]_3 - O(k_{\max}^2 e_1(Q)^2)) \geq E(Q) M^3,$$

using Lemma 3.1, and the number of ways to produce R using a switching of colour 11 is at most $k_{\max}^3 M_4$. Hence we can let $a_{11}(Q) = E(Q) M^3$ and $b_{11}(R) = k_{\max}^3 M_4$. This leads to

$$\alpha(Q, R) = \frac{k_{\max}^3 M_4}{E(Q) M^3} = O\left(\frac{k_{\max}^6}{M^2}\right) = o\left(\frac{k_{\max}^3}{M}\right).$$

3.10 Switchings of colour 12

Switchings of colour 12 reduce the number of non-loop edges with multiplicity 5 or 6, until this number is zero. They are applied to multigraphs Q such that switchings of colours $1, \dots, 11$ do not apply and $e_5(Q) + e_6(Q) \geq 1$.

Let Q be such a multigraph. Then a switching of colour 12 is defined by a sequence of distinct vertices

$$(v_0, w_0, v_1, w_1, v_2, w_2, v_3, w_3, v_4, w_4, v_5, w_5)$$

such that

- the edge $\{v_0, w_0\}$ has multiplicity D in Q , where $D \in \{5, 6\}$,
- each edge $\{v_j, w_j\}$ is a simple edge in Q , with multiplicity 1, for $j = 1, 2, 3, 4, 5$,

- the edges $\{v_0, v_j\}$ and $\{w_0, w_j\}$ all have multiplicity 0 in Q for $j = 1, 2, 3, 4, 5$ (that is, they are non-edges).

The switching reduces the multiplicity of the edge $\{v_0, w_0\}$ to $D - 5$, removes the edges $\{v_j, w_j\}$, $j = 1, 2, 3, 4, 5$, and inserts the simple edges $\{v_0, v_j\}$, $\{v_0, w_j\}$ for $j = 1, 2, 3, 4, 5$, as shown in Figure 9.




Figure 9: A switching of colour 12

Suppose that the multigraph R can be formed from Q by a switching of colour 12. The number of ways that a switching of colour 12 can be performed in Q is at least

$$2(e_5(Q) + e_6(Q)) ([2e_1(Q)]_5 - O(k_{\max}^2 e_1(Q)^4)) \geq (e_5(Q) + e_6(Q)) M^5,$$

using Lemma 3.1, and the number of ways that R can be produced by a switching of colour 12 is at most M_5^2 . Hence we can set $a_{12}(Q) = (e_5(Q) + e_6(Q))M^5$ and $b_{12}(R) = M_5^2$. This gives

$$\alpha(Q, R) = \frac{M_5^2}{(e_5(Q) + e_6(Q))M^5} = O\left(\frac{k_{\max}^8}{M^3}\right) = o\left(\frac{k_{\max}^3}{M}\right).$$

3.11 Switchings of colour 13

Switchings of colour 13 are used to reduce the number of simple loops until this number is at most $\lceil N_1/2 \rceil$. They are applied to multigraphs Q for which switchings of colours $1, \dots, 12$ do not apply and $\ell_1(Q) > \lceil N_1/2 \rceil$.

Let Q be such a multigraph. Then a switching of colour 13 is defined by a sequence of distinct vertices (v_0, v_1, v_2) such that there is a simple loop on v_0 in Q , the edge $\{v_1, v_2\}$ is a simple edge in Q , and the edges $\{v_0, v_1\}, \{v_0, v_2\}$ are both absent in Q (that is, they have multiplicity zero). The switching removes the simple loop at v_0 and the simple edge $\{v_1, v_2\}$ and inserts the two simple edges $\{v_0, v_1\}, \{v_0, v_2\}$, as shown in Figure 10.




Figure 10: A switching of colour 13

Suppose that the multigraph R can be obtained from Q by performing a switching of colour 13. The number of ways that a switching of colour 13 can be performed in Q is at least

$$\ell_1(Q)(2e_1(Q) - O(k_{\max}^2)) \geq \frac{1}{2}\ell_1(Q)M,$$

using Lemma 3.1, and the number of ways that a switching of colour 13 can produce R is at most M_2 . Therefore we can define $a_{13}(Q) = \frac{1}{2}\ell_1(Q)M$ and $b_{13}(R) = M_2$, leading to

$$\alpha(Q, R) = \frac{2M_2}{\ell_1(Q)M} \leq \frac{4M_2}{N_1M} \leq \frac{1}{120},$$

using the definition of N_1 .

3.12 Switchings of colour 14

Switchings of colour 14 reduce the number of non-loop edges with multiplicity 2 until this number is at most $\lceil N_2/2 \rceil$. This switching is applied to multigraphs Q such that switchings of colours $1, \dots, 13$ do not apply and $e_2(Q) > \lceil N_2/2 \rceil$.

Let Q be such a multigraph. A switching of colour 14 in Q is described by a sequence $(v_0, w_0, v_1, w_1, v_2, w_2)$ of distinct vertices such that

- $\{v_0, w_0\}$ is an edge of multiplicity 2 in Q ,
- $\{v_1, w_1\}$ and $\{v_2, w_2\}$ are simple edges in Q (with multiplicity 1),
- none of the edges $\{v_0, v_j\}$ or $\{w_0, w_j\}$ are present in Q , for $j = 1, 2$ (these are all non-edges, with multiplicity zero).

The switching removes these 3 edges and replaces them with two copies of $K_{1,2}$, one on $\{v_0, v_1, v_2\}$ centred at v_0 and the other on $\{w_0, w_1, w_2\}$ centred at w_0 . This operation is shown in Figure 11.




Figure 11: A switching of colour 14

Suppose that the multigraph R can be produced from Q using a switching of colour 14. The number of ways to perform a switching of colour 14 in Q is at least

$$2e_2(Q)([2e_1(Q)]_2 - O(k_{\max}^2 e_1(Q))) \geq e_2(Q)M^2,$$

using Lemma 3.1, and the number of ways that R can be produced using a switching of colour 14 is at most M_2^2 . Therefore we can take $a_{14}(Q) = e_2(Q)M^2$ and $b_{14}(R) = M_2^2$, leading to

$$\alpha(Q, R) = \frac{M_2^2}{e_2(Q)M^2} < \frac{2M_2^2}{N_2 M^2} \leq \frac{1}{120},$$

by the definition of N_2 .

3.13 Switchings of colour 15

Switchings of colour 15 are used to reduce the number of non-loop edges with multiplicity 3 until this number is at most $\lceil N_3/2 \rceil$. They are applied to multigraphs Q such that switchings of colours $1, \dots, 14$ do not apply and $e_3(Q) > \lceil N_3/2 \rceil$.

Let Q be such a multigraph. Then a switching of colour 15 is defined by a sequence of distinct vertices

$$(v_0, w_0, v_1, w_1, v_2, w_2, v_3, w_3)$$

such that

- $\{v_0, w_0\}$ has multiplicity 3 in Q ,
- each of $\{v_j, w_j\}$ is a simple edge in Q , for $j = 1, 2, 3$,
- the edges $\{v_0, v_j\}$ and $\{w_0, w_j\}$ are absent in Q for $j = 1, 2, 3$.

The switching removes the edges $\{v_j, w_j\}$ for $j = 0, 1, 2, 3$ (setting the multiplicity of each to zero) and inserts the simple edges $\{v_0, v_j\}, \{w_0, w_j\}$ for $j = 1, 2, 3$, as illustrated in Figure 8 for the case that $D = 3$.

Suppose that the multigraph R can be produced by performing a switching of colour 15 from Q . The number of ways to perform a switching of colour 15 in Q is at least

$$2e_3(Q) ([2e_1(Q)]_3 - O(k_{\max}^2 e_1(Q)^2)) \geq e_3(Q) M^3,$$

using Lemma 3.1, while the number of ways that R could be produced using a switching of colour 15 is at most M_3^2 . Therefore we can take $a_{15}(Q) = e_3(Q) M^3$ and $b_{15}(R) = M_3^2$, leading to

$$\alpha(Q, R) = \frac{M_3^2}{e_3(Q) M^3} \leq \frac{2 M_3^2}{N_3 M^3} \leq \frac{1}{120},$$

by the definition of N_3 .

3.14 Analysis of the switchings

We now explain how to apply Theorem 2.1 to analyse these switchings on $\mathcal{G}_0$. (See the statement of Theorem 2.1 for the necessary notation.)

We now define a directed graph Γ with vertex set $V(\Gamma) = \{v_Q : Q \in \mathcal{G}_0\}$, where v_Q is associated with the set $\mathcal{S}(v_Q) = \{Q\}$ containing one object. These sets are certainly disjoint. By a slight abuse of notation we will identify v_Q and Q from now on, and write Q for both. The edge set of Γ is defined as follows: there is an edge QR in Γ with colour $c \in \{1, \dots, 15\}$ if and only if R can be obtained from Q using a switching of colour c . Since a switching is used only where no switching with a lower-labelled colour applies, there is at most one edge from Q to R in Γ . Hence the endvertices of an edge uniquely determine the colour of the edge. We take $\lambda_c(v) = \frac{1}{15}$ for all c, v , and so for each edge QR in Γ we have

$$\hat{\alpha}(QR) = \frac{\alpha(QR)}{\lambda_{c(QR)}(R)} = 15 \alpha(QR).$$

Let Y, Z be as defined in (3.2), (3.3) and let $X = \mathcal{G}_0$.

Lemma 3.2. *With notation as established above, we have*

$$G(\mathbf{k}, J, J^*) = (1 + O(k_{\max}^3/M)) |\mathcal{G}_0 - Y|.$$

Proof. It follows from the analysis of the previous sections that $\alpha(QR) \leq \frac{1}{30}$, and therefore $\hat{\alpha}(QR) \leq \frac{1}{2}$, for all edges QR in Γ . Moreover, we have provided at least one switching which can be performed from Q , for each graph Q in $\mathcal{G}_0 - Z$. Hence X, Y, Z satisfy the requirements of Theorem 2.1 and we can conclude that

$$|Y| \leq \frac{\hat{\alpha}(Y, Z)}{1 - \hat{\alpha}(Y, Y)} |\mathcal{G}_0|,$$

where

$$\hat{\alpha}(W, W') = \max_{P \in \mathcal{P}(W, W')} \hat{\alpha}(P)$$

for all $W, W' \subseteq \mathcal{G}_0$.

Since $\mathcal{P}(Y, Y)$ by definition has only non-trivial paths, and $\hat{\alpha}(QR) \leq \frac{1}{2}$ for all edges in Γ , we know that $\hat{\alpha}(Y, Y) \leq \frac{1}{2}$, and therefore

$$|Y| \leq 2 \hat{\alpha}(Y, Z) |\mathcal{G}_0|.$$

Now let P be a path in Γ from some $Q \in Y$ to some element of Z , such that all internal vertices of P belong to $\mathcal{G}_0 - (Y \cup Z)$. Let QR be the first edge in P . Then QR cannot have colour in $\{1, \dots, 8\}$, since these switchings only produce graphs in Y . If QR is coloured with a colour in $\{9, 10, 11, 12\}$ then $\hat{\alpha}(QR) = O(k_{\max}^3/M)$, so $\hat{\alpha}(P) = O(k_{\max}^3/M)$.

The remaining possibility is that QR has colour $12+r$ for some $r \in \{1, 2, 3\}$, in which case P must contain at least $\lfloor N_r/2 \rfloor$ edges of colour $12+r$. Our analysis showed that $\alpha(e) \leq \frac{1}{120}$ and hence that $\hat{\alpha}(e) \leq \frac{1}{8}$, for all such edges e . By definition $\lfloor N_r/2 \rfloor \geq \frac{1}{2}(\log M - 1)$ for $r \in \{1, 2, 3\}$. Therefore,

$$\hat{\alpha}(P) \leq 8^{-(\log M - 1)/2} = O(M^{-1}),$$

which implies that $\hat{\alpha}(Y, Z) = O(k_{\max}^3/M)$. We conclude that

$$|\mathcal{G}_0 - Y| = (1 + O(k_{\max}^3/M)) |\mathcal{G}_0|.$$

But $\mathcal{G}_0 - Y \subseteq \mathcal{G}(\mathbf{k}, J, J^*) \subseteq \mathcal{G}_0$, and hence

$$G(\mathbf{k}, J, J^*) = (1 + O(k_{\max}^3/M)) |\mathcal{G}_0 - Y|,$$

completing the proof. $\square$

It remains to obtain an asymptotic expression for $\mathcal{G}_0 - Y$, which we do in the next section.

4 From pairings to multigraphs

In this section we work in the *pairing model* (also called *configuration model*), which we now describe. This model is standard for working with random graphs of fixed degrees: see for example [7]. Consider a set of M points partitioned into *cells* $c_1, \dots, c_n$, where cell c_i contains k_i points for $i = 1, 2, \dots, n$. Take a partition P (called a *pairing*) of the M points into $M/2$ pairs with each pair having the form $\{y, z\}$ where $y \in c_i$ and $z \in c_j$ for some i, j . The set of all such pairings, of which there are $M!/(M/2)! 2^{M/2}$, will be denoted by $\mathcal{C}(\mathbf{k})$.

A *loop* is a pair whose two points lie in the same cell, while a *link* is a pair involving two distinct cells. Two pairs are *parallel* if they involve the same cells. A *parallel class* is a maximal set of mutually parallel pairs. The *multiplicity* of a parallel class (and of the pairs in the class) is the cardinality of the class. As important special cases, a *simple pair* is a parallel class of multiplicity one, a *double pair* is a parallel class of multiplicity two and a *triple pair* is a parallel class of multiplicity three.

Each pairing gives rise to a multigraph in $\mathcal{G}(\mathbf{k})$ by replacing each cell by a vertex, and letting the multiplicity of the edge $\{v, w\}$ equal the multiplicity of the parallel class between the corresponding cells.

Let $\mathcal{C}_{\ell, d, t}$ be the set of all pairings in $\mathcal{C}(\mathbf{k})$ with exactly ℓ simple loops, exactly d double pairs and exactly t triple pairs, but with no loops of multiplicity greater than one and no links of multiplicity greater than three. If $G \in \mathcal{G}_0 - Y$ can be formed from a pairing $P \in \mathcal{C}_{\ell, d, t}$, then exactly

$$2^{-(\ell+d)} 6^{-t} \prod_{i=1}^n k_i!$$

pairings in $\mathcal{C}(\mathbf{k})$ give rise to G . Now defining

$$w(\ell, d, t) = 2^{\ell+d} 6^t |\mathcal{C}_{\ell, d, t}|,$$

we can write

$$|\mathcal{G}_0 - Y| = \left(\prod_{i=1}^n k_i! \right)^{-1} \sum_{\ell=0}^{N_1} \sum_{d=0}^{N_2} \sum_{t=0}^{N_3} w(\ell, d, t). \quad (4.1)$$

Hence it suffices to obtain an asymptotic expression for the above sum.

We will need the following two summation lemmas proved in [5]:

Lemma 4.1 ([5, Corollary 4.3]). *Let $0 \leq A_1 \leq A_2$ and $B_1 \leq B_2$ be real numbers. Suppose that there exist integers N, K with $N \geq 2$ and $0 \leq K \leq N$, and a real number $c > 2e$ such that $0 \leq Ac < N - K + 1$ and $|BN| < 1$ for all $A \in [A_1, A_2]$ and $B \in [B_1, B_2]$. Further suppose that there are real numbers δ_i , for $1 \leq i \leq N$, and $\gamma_i \geq 0$, for $0 \leq i \leq K$, such that*

$\sum_{j=1}^i |\delta_j| \leq \sum_{j=0}^K \gamma_j [i]_j < \frac{1}{5}$ for $1 \leq i \leq N$.

Given $A(1), \dots, A(N) \in [A_1, A_2]$ and $B(1), \dots, B(N) \in [B_1, B_2]$, define $n_0, n_1, \dots, n_N$ by $n_0 = 1$ and

$$\frac{n_i}{n_{i-1}} = \frac{A(i)}{i} (1 - (i-1)B(i)) (1 + \delta_i)$$

for $1 \leq i \leq N$, with the following interpretation: if $A(i) = 0$ then $n_j = 0$ for $i \leq j \leq N$. Then

$$\Sigma_1 \leq \sum_{i=0}^N n_i \leq \Sigma_2,$$

where

$$\begin{aligned} \Sigma_1 &= \exp\left(A_1 - \frac{1}{2}A_1^2 B_2 - 4 \sum_{j=0}^K \gamma_j (3A_1)^j\right) - \frac{1}{4}(2e/c)^N, \\ \Sigma_2 &= \exp\left(A_2 - \frac{1}{2}A_2^2 B_1 + \frac{1}{2}A_2^3 B_1^2 + 4 \sum_{j=0}^K \gamma_j (3A_2)^j\right) + \frac{1}{4}(2e/c)^N. \quad \square \end{aligned}$$

Lemma 4.2 ([5, Corollary 4.5]). Let $N \geq 2$ be an integer and, for $1 \leq i \leq N$, let real numbers $A(i)$, $B(i)$ be given such that $A(i) \geq 0$ and $1 - (i-1)B(i) \geq 0$. Define $A_1 = \min_{i=1}^N A(i)$, $A_2 = \max_{i=1}^N A(i)$, $C_1 = \min_{i=1}^N A(i)B(i)$ and $C_2 = \max_{i=1}^N A(i)B(i)$. Suppose that there exists a real number $\hat{c}$ with $0 < \hat{c} < \frac{1}{3}$ such that $\max\{A/N, |C|\} \leq \hat{c}$ for all $A \in [A_1, A_2]$, $C \in [C_1, C_2]$. Define $n_0, \dots, n_N$ by $n_0 = 1$ and

$$\frac{n_i}{n_{i-1}} = \frac{A(i)}{i} (1 - (i-1)B(i))$$

for $1 \leq i \leq N$, with the following interpretation: if $A(i) = 0$ or $1 - (i-1)B(i) = 0$, then $n_j = 0$ for $i \leq j \leq N$. Then

$$\Sigma_1 \leq \sum_{i=0}^N n_i \leq \Sigma_2,$$

where

$$\begin{aligned} \Sigma_1 &= \exp\left(A_1 - \frac{1}{2}A_1 C_2\right) - (2e\hat{c})^N, \\ \Sigma_2 &= \exp\left(A_2 - \frac{1}{2}A_2 C_1 + \frac{1}{2}A_2 C_1^2\right) + (2e\hat{c})^N. \quad \square \end{aligned}$$

First we perform a summation over the number of edges of multiplicity three.

Lemma 4.3. Uniformly for $0 \leq d \leq N_2$ and $0 \leq \ell \leq N_1$, we have

$$\sum_{t=0}^{N_3} \frac{w(\ell, d, t)}{w(\ell, d, 0)} = \exp\left(\frac{M_3^2}{2M^3} + O(k_{\max}^3/M)\right).$$

Proof. We will apply Lemma 4.2. Let t' be the first value of $t \leq N_3$ for which $\mathcal{C}_{\ell,d,t} = \emptyset$, or $t' = N_3 + 1$ if there is no such value. By [12, Lemma 4.1] we have

$$\frac{w(\ell, d, t)}{w(\ell, d, t-1)} = \frac{6|\mathcal{C}_{\ell,d,t}|}{|\mathcal{C}_{\ell,d,t-1}|} = \frac{M_3^2 - \alpha_t k_{\max}^2 (k_{\max}^2 + \ell + d + t - 1) M_3}{2t M^3}, \quad (4.2)$$

where $\alpha_t = \alpha_t(\ell, d)$ is bounded independently of ℓ, d and M , for $1 \leq t < t'$. Note that our expression for N_3 is no smaller than, and at most a constant factor larger than, that used in [12]. Examination of the proof of [12, Lemma 4.1] shows that the expression given there for $|\mathcal{C}_{\ell,d,t}|/|\mathcal{C}_{\ell,d,t-1}|$ also holds for $1 \leq t \leq N_3$.

For $1 \leq t < t'$, define

$$A(t) = \frac{M_3^2 - \alpha_t k_{\max}^2 (k_{\max}^2 + \ell + d) M_3}{2M^3}, \quad C(t) = \frac{\alpha_t k_{\max}^2 M_3}{2M^3}.$$

Now $M_3 > 0$ since $1 \leq t < t'$. If $\alpha_t \leq 0$ then by definition $A(t) \geq M_3^2/2M^3 > 0$. Now suppose that $\alpha_t > 0$. Then $C(t) > 0$. Note that the left hand side of (4.2) is positive, and multiplying by t we see that $A(t) - (t-1)C(t)$ is positive, which implies that $A(t) > 0$. Therefore $A(t) > 0$ whenever $1 \leq t < t'$. Define $B(t) = C(t)/A(t)$ for $1 \leq t < t'$. Also define $A(t) = B(t) = 0$ for $t' \leq t \leq N_3$.

Define A_1, A_2, C_1, C_2 by taking the minimum and maximum of $A(t)$ and $C(t)$ over $1 \leq t \leq N_3$, as in Lemma 4.2. Let $A \in [A_1, A_2]$ and $C \in [C_1, C_2]$, and set $\hat{c} = \frac{1}{80}$. Since $A = M_3^2/2M^3 + o(1)$ and $C = o(1)$, we have that $\max\{A/N_3, |C|\} < \hat{c}$ for M sufficiently large, by the definition of N_3 .

Therefore Lemma 4.2 applies and says that

$$\sum_{t=0}^{N_3} \frac{w(\ell, d, t)}{w(\ell, d, 0)} = \exp\left(\frac{M_3^2}{2M^3} + O(k_{\max}^4 (k_{\max}^2 + \ell + d)/M^2)\right) + O((e/40)^{N_3}).$$

Since $\ell + d \leq N_1 + N_2 = O(k_{\max}^2 + \log M)$, the relative error term is $O(k_{\max}^3/M)$, as claimed. Finally, $(e/40)^{N_3} \leq (e/40)^{\log M} \leq M^{-2}$. Since the sum we are estimating is at least equal to one, the additive error term is covered by the error term inside the exponential. This completes the proof. $\square$

Next we perform a summation over the number of simple loops.

Lemma 4.4. *Uniformly for $0 \leq d \leq N_2$, we have*

$$\sum_{\ell=0}^{N_1} \frac{w(\ell, d, 0)}{w(0, d, 0)} = \exp\left(\frac{M_2}{M} + O\left(\frac{k_{\max} d}{M} + \frac{k_{\max}^3}{M}\right)\right).$$

Proof. Let ℓ' be the first value of $\ell \leq N_1$ for which $\mathcal{C}_{\ell,d,0} = \emptyset$, or $\ell' = N_1 + 1$ if there is no such value. By [12, Lemma 4.2], for $1 \leq \ell < \ell'$ we have

$$\frac{w(\ell, d, 0)}{w(\ell-1, d, 0)} = \frac{2 |\mathcal{C}_{\ell,d,0}|}{|\mathcal{C}_{\ell-1,d,0}|} = \frac{M_2 - \beta_\ell(k_{\max}^3 + k_{\max}d + k_{\max}^2(\ell-1))}{\ell M}, \quad (4.3)$$

where $\beta_\ell = \beta_\ell(d)$ is uniformly bounded (independently of d and M). As in Lemma 4.3 above, our expression for N_1 is no smaller than, and at most a constant factor larger than, that used in [12]. Examination of the proof of [12, Lemma 4.2] shows that the expression given there for $|\mathcal{C}_{\ell,d,0}|/|\mathcal{C}_{\ell-1,d,0}|$ also holds for $1 \leq \ell \leq N_1$. The proof is completed by adapting the argument used for Lemma 4.3, above.

For $1 \leq \ell < \ell'$, define

$$A(\ell) = \frac{M_2 - \beta_\ell(k_{\max}^3 + k_{\max}d)}{M}, \quad C(\ell) = \frac{\beta_\ell k_{\max}^2}{M}.$$

By the same argument as in the previous lemma, $A(\ell) > 0$ for $\ell < \ell'$. Define $B(\ell) = C(\ell)/A(\ell)$ for $1 \leq \ell < \ell'$. Also define $A(\ell) = B(\ell) = 0$ for $\ell' \leq \ell \leq N_1$, and $\hat{c} = \frac{1}{80}$. Lemma 4.2 now applies to give the desired result. $\square$

Next we perform a summation over the number of edges of multiplicity two. The exponential factor in the summand corresponds to the error term from Lemma 4.4 which depends on d .

Lemma 4.5. *For any constant ρ have*

$$\sum_{d=0}^{N_2} \frac{w(0, d, 0)}{w(0, 0, 0)} \exp\left(\frac{\rho k_{\max} d}{M}\right) = \exp\left(\frac{M_2^2}{2M^2} - \frac{M_3^2}{2M^3} + O(k_{\max}^3/M)\right).$$

Proof. We divide the proof into two cases, following the division used in [12, Lemmas 4.3–4.4]. Again, our expression for N_2 is no smaller than, and at most a constant factor larger than, that used in [12]. Examination of the proofs of [12, Lemmas 4.3–4.4] shows that the expression given there for $|\mathcal{C}_{0,d,0}|/|\mathcal{C}_{0,d-1,0}|$ also holds for $1 \leq d \leq N_2$.

For $0 \leq d \leq N_2$, define

$$m_d = \frac{w(0, d, 0)}{w(0, 0, 0)} \exp\left(\frac{\rho k_{\max} d}{M}\right).$$

First suppose that $M_2 \leq M$. From [12, Lemma 4.3] we know that for $1 \leq d \leq N_2$ with $\mathcal{C}_{0,d,0} \neq \emptyset$,

$$\frac{m_d}{m_{d-1}} = \frac{2 \exp(\rho k_{\max}/M) |\mathcal{C}_{0,d,0}|}{|\mathcal{C}_{0,d-1,0}|} = \frac{M_2^2}{2dM^2} + O\left(\frac{k_{\max}(k_{\max}^2 + d)M_2}{dM^2}\right).$$

The current lemma follows from this, arguing as in the proofs of the previous two lemmas. (Note that the term $M_3^2/2M^3$ in the answer is absorbed into the error term, by the assumption that $M_2 \leq M$.)

Now suppose that $M_2 > M$. By [12, Lemma 4.4] we have for $1 \leq d \leq N_2$ with $\mathcal{C}_{0,d,0} \neq \emptyset$,

$$\frac{m_d}{m_{d-1}} = \frac{2 \exp(\rho k_{\max}/M) |\mathcal{C}_{0,d,0}|}{|\mathcal{C}_{0,d-1,0}|} = \frac{A(d)}{d} (1 - (d-1)B) (1 + \delta_d),$$

where

$$\begin{aligned} A(d) &= \frac{M_2^2}{2M^2} + \frac{2M_2^2 M_3}{M^4} - \frac{M_3^2}{2M^3} - \frac{M_2^4}{M^5} + O(k_{\max}^3/M), \\ B &= -\frac{8}{M} + \frac{16M_3}{M_2^2}, \\ \delta_d &= O((d-1)/M_2) \quad \text{uniformly for } 1 \leq d \leq N_2. \end{aligned}$$

(Note that $A(d) \exp(\rho k_{\max}/M)$ equals $A(d)$ within the precision afforded by the error term, since $k_{\max} A(d)/M = O(k_{\max} M_2^2/M^3) = O(k_{\max}^3/M)$.) Again, our N_2 is no smaller than, and at most a constant factor larger than, the value of N_2 used in [12], and examination of the proof of [12, Lemma 4.4] shows that the expression for $|\mathcal{C}_{0,d,0}|/|\mathcal{C}_{0,d-1,0}|$ also holds for $1 \leq d \leq N_2$.

Let d' be the first value of $d \leq N_2$ for which $\mathcal{C}_{0,d,0} = \emptyset$, or $d' = N_2 + 1$ if no such value of d exists. Define $n_d = m_d/m_{d-1}$ for $0 \leq d < d'$ and $n_d = 0$ for $d \geq d'$. We will apply Lemma 4.1 to $\sum_d n_d$.

Let $A_1 = \min_d A(d)$ and $A_2 = \max_d A(d)$, with the minimum and maximum taken over $1 \leq d \leq N_2$, and let $B_1 = B_2 = B$. Define $c = 80$ and $K = 2$. Then the conditions of Lemma 4.1 apply with $\gamma_0 = \gamma_1 = 0$ and some value of γ_2 which satisfies $\gamma_2 = O(1/M_2)$. Application of that lemma now gives the desired result. $\square$

Recall the definition of $y_1, x_2, x_3 \in \{0, 1\}$ given just before the statement of Theorem 1.1. By combining the last three summations we obtain the following.

Lemma 4.6. *As $M \rightarrow \infty$ and $k_{\max}^3 = o(M)$, we have*

$$\sum_{d=0}^{x_2 N_2} \sum_{\ell=0}^{y_1 N_1} \sum_{t=0}^{x_3 N_3} \frac{w(\ell, d, t)}{w(0, 0, 0)} = \exp \left(y_1 \frac{M_2}{M} + x_2 \frac{M_2^2}{2M^2} + (x_3 - x_2) \frac{M_3^2}{2M^3} + O(k_{\max}^3/M) \right).$$

Proof. From Lemmas 4.3 and 4.4, we have

$$\sum_{\ell=0}^{y_1 N_1} \sum_{t=0}^{x_3 N_3} \frac{w(\ell, d, t)}{w(0, 0, 0)} = \frac{w(0, d, 0)}{w(0, 0, 0)} \exp \left(y_1 \frac{M_2}{M} + x_3 \frac{M_3^2}{2M^3} + O \left(\frac{k_{\max} d}{M} + \frac{k_{\max}^3}{M} \right) \right) \quad (4.4)$$

uniformly over d .

If $x_2 = 0$, we are finished. If $x_2 = 1$, choose two constants ρ_1, ρ_2 such that the actual value of the error term $O(k_{\max}d/M)$ in (4.4) lies in $[\rho_1 k_{\max}d/M, \rho_2 k_{\max}d/M]$ for all $d \in \{1, \dots, N_2\}$. Applying Lemma 4.5 with $\rho \in \{\rho_1, \rho_2\}$ and noting that the result does not depend on ρ within the precision given by the error term, we are done. $\square$

Finally we can prove our main result.

Proof of Theorem 1.1. From Lemma 3.2 and Lemma 4.6 together with (4.1), we obtain

$$\begin{aligned} G(\mathbf{k}, J, J^*) &= (1 + O(k_{\max}^3/M)) |\mathcal{G}_0 - Y| \\ &= \frac{1 + O(k_{\max}^3/M)}{k_1! \dots k_n!} \sum_{d=0}^{x_2 N_2} \sum_{\ell=0}^{y_1 N_2} \sum_{t=0}^{x_3 N_3} w(\ell, d, t) \\ &= \frac{w(0, 0, 0)}{k_1! \dots k_n!} \exp\left(y_1 \frac{M_2}{M} + x_2 \frac{M_2^2}{2M^2} + (x_3 - x_2) \frac{M_3^2}{2M^3} + O(k_{\max}^3/M)\right). \end{aligned}$$

Now $w(0, 0, 0) = |\mathcal{C}_{0,0,0}|$, and it follows from [12, Lemma 5.1] that

$$|\mathcal{C}_{0,0,0}| = \frac{M!}{(M/2)! 2^{M/2}} \exp\left(-\frac{M_2}{2M} - \frac{M_2^2}{4M^2} - \frac{M_2^2 M_3}{2M^4} + \frac{M_2^4}{4M^5} + \frac{M_3^2}{6M^3} + O(k_{\max}^3/M)\right).$$

Combining these two expressions completes the proof. $\square$

5 Comparison to a naïve model

Let $p \in (0, 1)$ be a probability which we will define later. Abusing notation slightly, we define the functions

$$J(z) = \sum_{j \in J} p^j z^j, \quad J^*(z) = \sum_{j \in J^*} p^j z^j$$

and the probability generating functions

$$f(z) = \frac{J(z)}{J(1)}, \quad g(z) = \frac{J^*(z)}{J^*(1)}.$$

Consider a random symmetric nonnegative integer matrix $A = (a_{ij})$ which is created as follows:

- All entries on and above the diagonal are independent;
- For $i = 1, \dots, n$, let a_{ii} be a randomly chosen element of J^* , with $\Pr(a_{ii} = b) = p^b / J^*(1)$ for all $b \in J^*$;

- For $1 \leq i < j \leq n$ let a_{ij} be a randomly chosen element of J , with $\Pr(a_{ij} = b) = p^b/J(1)$ for all $b \in J$;
- Finally, let $a_{ji} = a_{ij}$ for $1 \leq i < j \leq n$.

We write $\Pr(\cdot)$ to mean probabilities generated by the above procedure. If A_0 is a fixed matrix which corresponds to an element of $\mathcal{G}(\mathbf{k}, J, J^*)$ then the probability that A_0 is produced by the above procedure is

$$\Pr(A_0) = \frac{p^{M/2}}{J(1)^{\binom{n}{2}} J^*(1)^n}. \quad (5.1)$$

This follows as the probability of A_0 depends only on the sum of the diagonal and above-diagonal entries (that is, on the number of edges of the corresponding multigraph).

For $i = 1, \dots, n$, let $\mathcal{E}_i$ denote the event that the sum of row i equals k_i , recalling that diagonal entries are weighted by a factor of 2 as in (1.1). Now

$$G(\mathbf{k}, J, J^*) = \frac{\Pr(\mathcal{E}_1 \wedge \mathcal{E}_2 \wedge \dots \wedge \mathcal{E}_n)}{\Pr(A_0)}$$

since $\Pr(\cdot)$ is uniform on (the set of matrices corresponding to) $\mathcal{G}(\mathbf{k}, J, J^*)$. By (incorrectly) assuming that the events $\mathcal{E}_1, \dots, \mathcal{E}_n$ are independent, we obtain the following naïve estimate of $G(\mathbf{k}, J, J^*)$, using (5.1):

$$\begin{aligned} G_p(\mathbf{k}, J, J^*) &= \frac{\prod_{i=1}^n \Pr(\mathcal{E}_i)}{\Pr(A_0)} \\ &= \frac{J(1)^{\binom{n}{2}} J^*(1)^n}{p^{M/2}} \prod_{i=1}^n [z^{k_i}] f(z)^{n-1} g(z^2) \\ &= p^{-M/2} J(1)^{-\binom{n}{2}} \prod_{i=1}^n [z^{k_i}] (J(z)^{n-1} J^*(z^2)). \end{aligned} \quad (5.2)$$

Let $k = M/n$ denote the target average row sum. The value p_0 of p which makes the expected row sum equal to k satisfies

$$p_0 = \frac{k}{n} + \frac{(1 - 2x_2)k^2}{n^2} + O(k/n^2). \quad (5.3)$$

We write $G_{\text{naïve}}(\mathbf{k}, J, J^*)$ for $G_{p_0}(\mathbf{k}, J, J^*)$ from now on. (As we shall see, the exact value of the $O(k/n^2)$ term does not affect Theorem 5.1 below, within the stated error bound, even though some of the intermediate formulae we give in the proof are affected.)

We now compare the expression of Theorem 1.1 with the naïve estimate $G_{\text{naïve}}(\mathbf{k}, J, J^*)$. For future applications, it will be convenient to express the result in terms of the scaled

central moments μ_2, μ_3 defined by

$$\mu_r = \frac{1}{M} \sum_{i=1}^n (k_i - k)^r$$

for $r = 2, 3$.

Theorem 5.1. *Under the conditions of Theorem 1.1, we have*

$$\begin{aligned} G(\mathbf{k}, J, J^*) &= \sqrt{2} G_{\text{naive}}(\mathbf{k}, J, J^*) \exp \left(\frac{1}{4} (1 - \mu_2) (1 + 2x_2 + \mu_2(1 - 2x_2)) \right. \\ &\quad + \frac{6\mu_2\mu_3(x_3 - x_2) - \mu_2^3}{2n} + \frac{3\mu_2^2(\mu_2^2 - 2\mu_3) + 2\mu_3^2(3x_3 - 3x_2 + 1)}{12M} \\ &\quad \left. + \frac{\mu_2^2 M}{2n^2} (9x_3 - 9x_2 - 1) + O(k_{\max}^3/M) \right). \end{aligned} \quad (5.4)$$

Proof. Suppose that

$$p = \frac{k}{n} + \frac{(1 - 2x_2)k^2}{n^2} + \frac{ck}{n^2},$$

where $c = c(n) = O(1)$. By direct computation we find that

$$p^{M/2} = \left(\frac{k}{n} \right)^{M/2} \exp \left(\frac{1}{2} k^2 (1 - 2x_2) - \frac{k^3}{4n} + \frac{ck}{2} + O(k_{\max}^3/M) \right) \quad (5.5)$$

and

$$\begin{aligned} J(1)^{\binom{n}{2}} &= \exp \left(\binom{n}{2} p - \frac{(1 - 2x_2)p^2 n^2}{4} + \frac{(1 - 3x_2 + 3x_3)p^3 n^2}{6} + O(k_{\max}^3/M) \right) \\ &= \exp \left(\frac{1}{2} M - \frac{1}{4} k(2x_2 k + 2 - k) + \frac{ck}{2} - \frac{k^3(2 + 3x_2 - 3x_3)}{6n} + O(k_{\max}^3/M) \right). \end{aligned} \quad (5.6)$$

To calculate the product in (5.2) we consider two cases, depending on the value of x_2 . First assume that $x_2 = 0$ (that is, the value 2 is not permitted off the diagonal). Then

$$J(z)^{n-1} J^*(z^2) = (1 + pz)^{n-1} H(z),$$

where

$$H(z) = \left(1 + \frac{\sum_{i \geq 3} x_i p^i z^i}{1 + pz} \right)^{n-1} \left(1 + \sum_{i \geq 1} y_i p^i z^{2i} \right).$$

Consequently, for any integer r with $0 \leq r \leq k_{\max}$ we have

$$\begin{aligned} [z^r] J(z)^{n-1} J^*(z^2) &= \sum_{j=0}^r \binom{n-1}{r-j} p^{r-j} [z^j] H(z) \\ &= \binom{n-1}{r} p^r + \binom{n-1}{r-2} y_1 p^{r-1} + \binom{n-1}{r-3} x_3 (n-1) p^r + \binom{n-1}{r} p^r \Delta, \end{aligned} \quad (5.7)$$

where

$$\Delta = \sum_{j=4}^r \binom{n-1}{r-j} \binom{n-1}{r}^{-1} p^{-j} [z^j] H(z).$$

Now define

$$H^+(z) = \left(1 + \frac{\sum_{i \geq 3} p^i z^i}{1 - pz}\right)^{n-1} \left(1 + \sum_{i \geq 1} p^i z^{2i}\right) = \left(1 + \frac{p^3 z^3}{(1 - pz)^2}\right)^{n-1} (1 - pz^2)^{-1}.$$

Notice that the coefficients of $H^+(z)$, when expanded as a Taylor series in z , are nonnegative and dominate those of $H(z)$. Also notice that $\binom{n-1}{r-j} \binom{n-1}{r}^{-1} = O(1)(r/n)^j$ uniformly for $4 \leq j \leq r$, since $r = o(n^{1/2})$. Therefore, setting $\alpha = r/pn$,

$$\begin{aligned} |\Delta| &= O(1) \sum_{j=4}^r \alpha^j [z^j] H^+(z) \\ &= O(H^+(\alpha) - 1 - p\alpha^2 - (n-1)p^3\alpha^3) \\ &= O(rk_{\max}^3/M^2), \end{aligned}$$

where the last line follows since $H^+(z) - 1 - pz^2 - (n-1)p^3z^3$ has three zero derivatives at $z = 0$ and its fourth derivative is $O(p^2)$ for $z = O(k_{\max})$. Applying this bound to (5.7), we have

$$\begin{aligned} [z^r] J(z)^{n-1} J^*(z^2) &= \binom{n-1}{r} p^r \exp\left(\frac{y_1 r(r-1)}{kn} + \frac{x_3 r^3}{n^2} + O(rk_{\max}^3/M^2)\right) \\ &= \frac{n^r p^r}{r!} \exp\left(-\frac{r(r+1)}{2n} + \frac{y_1 r(r-1)}{kn} + \frac{r^3(6x_3-1)}{6n^2} + O(rk_{\max}^3/M^2)\right). \end{aligned} \quad (5.8)$$

Now we assume that $x_2 = 1$ and repeat the calculation. We have

$$J(z)^{n-1} J^*(z^2) = (1 - pz)^{-n+1} \left((1 - pz) \sum_{i \geq 0} x_i p^i z^i \right)^{n-1} \left(1 + \sum_{i \geq 1} y_i p^i z^{2i} \right).$$

Arguing as before, we find that for any r with $0 \leq r \leq k_{\max}$,

$$\begin{aligned} [z^r] J(z)^{n-1} J^*(z^2) &= \binom{n+r-2}{r} p^r + \binom{n+r-4}{r-2} y_1 p^{r-1} + \binom{n+r-5}{r-3} (x_3-1)(n-1) p^r \\ &\quad + \binom{n+r-2}{r} p^r O(rk_{\max}^3/M^2) \\ &= \binom{n+r-2}{r} p^r \exp\left(\frac{y_1 r(r-1)}{kn} + \frac{(x_3-1)r^3}{n^2} + O(rk_{\max}^3/M^2)\right) \\ &= \frac{n^r p^r}{r!} \exp\left(\frac{r(r-3)}{2n} + \frac{y_1 r(r-1)}{kn} + \frac{r^3(6x_3-7)}{6n^2} + O(rk_{\max}^3/M^2)\right). \end{aligned} \quad (5.9)$$

Combining the cases (5.8) and (5.9) we have, for $i = 1, \dots, n$,

$$[z^{k_i}] J(z)^{n-1} J^*(z^2) = \frac{n^{k_i} p^{k_i}}{k_i!} \exp \left(- \left(\frac{1+2x_2}{2n} + \frac{y_1}{kn} \right) k_i - \left(\frac{1-2x_2}{2n} - \frac{y_1}{kn} \right) k_i^2 - \frac{1+6x_2-6x_3}{6n^2} k_i^3 + O(k_i k_{\max}^3/M^2) \right).$$

Multiplying these n equations together gives

$$\begin{aligned} & \prod_{i=1}^n [z^{k_i}] J(z)^{n-1} J^*(z^2) \\ &= \frac{n^M p^M}{\prod_{i=1}^n k_i!} \exp \left(\frac{M_2(2x_2-1) - 2M}{2n} + \frac{y_1 M_2}{kn} - \frac{(6x_2-6x_3+1)M_3}{6n^2} + O(k_{\max}^3/M) \right). \end{aligned} \quad (5.10)$$

We now substitute (5.5), (5.6) and (5.10) into (5.2). Writing the result in terms of μ_2 , μ_3 , M and n , using the identities

$$\begin{aligned} M_2 &= M\mu_2 + (k-1)M, \\ M_3 &= M\mu_3 + 3(k-1)M\mu_2 + (k-1)(k-2)M \end{aligned}$$

we find that

$$\begin{aligned} & G_{\text{naive}}(\mathbf{k}, J, J^*) \\ &= \frac{1}{\prod_{i=1}^n k_i!} \left(\frac{M}{e} \right)^{M/2} \exp \left(-y_1(1-\mu_2) + \frac{(2y_1+2x_2(\mu_2-1)-\mu_2)k}{2} + \frac{(2x_2-1)k^2}{4} - \frac{k(6x_2-6x_3+1)(2\mu_3+6\mu_2k+k^2)}{12n} + O(k_{\max}^3/M) \right). \end{aligned}$$

(At this point we can observe that no remaining terms depend on c , which verifies our earlier claim that the exact value of the $O(k/n^2)$ term in (5.3) does not affect the statement of this theorem.) The proof is completed by applying Theorem 1.1 and using Stirling's formula. $\square$

Corollary 5.2. *Under the conditions of Theorem 5.1, if $\mu_2 = O(M^{1/6})$ then*

$$\begin{aligned} & G(\mathbf{k}, J, J^*) \\ &= \sqrt{2} G_{\text{naive}}(\mathbf{k}, J, J^*) \exp \left(\frac{1}{4} (1-\mu_2)(1+2x_2+\mu_2(1-2x_2)) + O(k_{\max}^2/M^{2/3}) \right). \end{aligned}$$

If the stronger bound $\mu_2 = O(k_{\max}^{1/2})$ holds then

$$\begin{aligned} & G(\mathbf{k}, J, J^*) \\ &= \sqrt{2} G_{\text{naive}}(\mathbf{k}, J, J^*) \exp \left(\frac{1}{4} (1-\mu_2)(1+2x_2+\mu_2(1-2x_2)) + O(k_{\max}^3/M) \right). \end{aligned}$$

Finally, if $\mathbf{k} = (k, k, k, \dots, k)$ is a regular degree sequence with kn even, then

$$G(\mathbf{k}, J, J^*) = \sqrt{2} \exp\left(\frac{1}{4}(1 + 2x_2) + O(k^2/n)\right) G_{\text{naive}}(\mathbf{k}, J, J^*). \quad (5.11)$$

Proof. For the first two statements, we just need to check that all additional terms inside the exponential factor in Theorem 5.1 are covered by the claimed error bounds. For this it is useful to note that $\mu_2 \leq k_{\max}$ and $|\mu_3| \leq k_{\max}\mu_2$. Finally, when $\mathbf{k}$ is a regular degree sequence we have $\mu_2 = \mu_3 = 0$ and (5.11) follows. $\square$

The constant $\frac{1}{4}(1 + 2x_2)$ in (5.11) was previously noted for simple sparse regular graphs in [12], and in [4] for simple sparse regular graphs with loops allowed. Interestingly, with different error terms, the same constant was observed in [12] for dense simple regular graphs, in [4] for dense simple regular graphs with loops allowed, and in [9] for dense symmetric integer matrices with zero diagonal. In these three cases, we conjectured that (5.11) holds (with some vanishing error term) for all degree sequences except for the two extreme cases of graphs with no edges and graphs with all possible edges. The corresponding conjectures may be less likely to be true in full generality for all J, J^* .

From the first expression of Corollary 5.2, we see that $G(\mathbf{k}, J, J^*)$ is closely approximated by $\sqrt{2} G_{\text{naive}}(\mathbf{k}, J, J^*)$ whenever μ_2 is close to 1. It appears that, in the random matrix model described at the start of this section, μ_2 will be concentrated around 1 with high probability whenever p tends to zero slowly enough to ensure that $M \rightarrow \infty$ with high probability, but quickly enough to ensure that $k_{\max}^3 = o(M)$ with high probability. If so, this will lead to a model for the degree sequences of sparse multigraphs analogous to that obtained by McKay and Wormald [13] for graphs, and by McKay and Skerman [10] for bipartite graphs and directed graphs. Details will be given in a future paper.

References

- [1] A. Barvinok and J. A. Hartigan, An asymptotic formula for the number of non-negative integer matrices with prescribed row and column sums, *Trans. Amer. Math. Soc.*, **364** (2012), 4323–4368.
- [2] E. A. Bender and E. R. Canfield, The asymptotic number of labeled graphs with given degree sequences, *J. Combin. Theory Ser. A*, **24** (1978) 296–307.
- [3] C. Greenhill and B. D. McKay, Asymptotic enumeration of sparse nonnegative integer matrices with specified row and column sums, *Adv. in Appl. Math.*, **41** (2008), 459–481.

- [4] C. Greenhill and B. D. McKay, Counting loopy graphs with given degrees, *Linear Alg. Appl.*, **436** (2012), 901–926.
- [5] C. Greenhill, B. D. McKay and X. Wang, Asymptotic enumeration of sparse 0-1 matrices with irregular row and column sums, *J. Combin. Theory Ser. A*, **113** (2006), 291–324.
- [6] M. Hasheminezhad and B. D. McKay, Combinatorial estimates by the switching method, *Contemp. Math.*, **531** (2010), 209–221.
- [7] B. D. McKay, Asymptotics for 0-1 matrices with prescribed line sums, in *Enumeration and Design*, Academic Press, Toronto, 1984, pp. 225–238.
- [8] B. D. McKay, Subgraphs of dense random graphs with specified degrees, *Combin. Probab. Comput.*, **20** (2011) 413–433.
- [9] B. D. McKay and J. C. McLeod, Asymptotic enumeration of symmetric integer matrices with uniform row sums, *J. Aust. Math. Soc.*, to appear (2013).
<http://arxiv.org/abs/1108.4496>
- [10] B. D. McKay and F. Skerman, Degree sequences of random digraphs and bipartite graphs, submitted (2012). <http://arxiv.org/abs/1302.2446>
- [11] B. D. McKay and N. C. Wormald, Asymptotic enumeration by degree sequence of graphs of high degree, *European J. Combin.*, **11** (1990) 565–580.
- [12] B. D. McKay and N. C. Wormald, Asymptotic enumeration by degree sequence of graphs with degrees $o(n^{1/2})$, *Combinatorica*, **11** (1991) 369–382.
- [13] B. D. McKay and N. C. Wormald, The degree sequence of a random graph. I. The models, *Random Structures and Algorithms* **11** (1997) 97–117.
- [14] R. C. Read, *Some enumeration problems in graph theory*, Doctoral Thesis, University of London (1958), p. 156.